

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number  
**WO 03/053342 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K**
- (21) International Application Number: **PCT/US02/39182**
- (22) International Filing Date: 9 December 2002 (09.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
10/006,430 10 December 2001 (10.12.2001) US
- (71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **GRAHAM, Mark, J.** [US/US]; 2305 S. Ola Vista, San Clemente, CA 92672 (US). **DOBIE, Kenneth, W.** [US/US]; 703 Stratford Court #4, Del Mar, CA 92014 (US).
- (74) Agents: **LICATA, Jane, Massey et al.**; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTISENSE MODULATION OF CD81 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of CD81. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding CD81. Methods of using these compounds for modulation of CD81 expression and for treatment of diseases associated with expression of CD81 are provided.

WO 03/053342 A2

## ANTISENSE MODULATION OF CD81 EXPRESSION

### FIELD OF THE INVENTION

The present invention provides compositions and methods for modulating the expression of CD81. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding CD81. Such compounds have been shown to modulate the expression of CD81.

### BACKGROUND OF THE INVENTION

The tetraspanins (also known as the TM4 superfamily or TM4SF) comprise a large family of proteins originally identified as leukocyte antigens. They have been shown to influence cell adhesion and migration, to alter cell morphology and to affect the activation state of a cell. Antibodies to tetraspanins induce intracellular signaling that is required for functional effects such as adhesion. A common characteristic of tetraspanins is a propensity to physically associate with a variety of other membrane proteins such as integrins, lineage specific molecules and other tetraspanins (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109).

CD81 (also known as target of antiproliferative antibody 1, TAPA-1 and CD81 antigen) is a 26-kDa cell surface protein composed of four transmembrane and two extracellular domains. It exhibits an overall topology common to the tetraspanins (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109). CD81 has been cloned and mapped to chromosome 11p15.5 (Oren et al., *Mol. Cell Biol.*, 1990, 10, 4007-4015; Virtaneva et al., *Immunogenetics*, 1994, 39, 329-334).

Disclosed and claimed in PCT publication WO 99/19198 are a transgenic non-human animal carrying a transgene encoding CD81

RTSP-0443

-2-

protein, a nucleic acid molecule which encodes a CD81 protein, and a nucleic acid molecule which hybridizes to said nucleic acid molecule under both standard conditions and conditions of high stringency (Abrignani and Grandi, 1999).

CD81 is expressed on the cell surfaces of most human tissues with the notable exceptions of red blood cells and platelets. The fact that CD81 was discovered in many different biological assays implies that it is involved in many different cellular functions that are manifested differently in the various cell types (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109).

CD81 forms several different molecular complexes. It associates on the surface of B-cells with B-cell-specific molecules CD19 and CD21, interferon inducible antigen Leu-13, MHC-class II molecules, integrins and other tetraspanins (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109). CD81 associations on the surface of T-cells include CD82, CD4, CD8 and alpha-4-beta-1 integrin (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109). Cross-linking of CD81 to various cell types may affect different sets of associated proteins expressed by the cells.

Biological functions of CD81 have been delineated via the use of anti-CD81 antibodies and include antiproliferative effects, adhesion, induction of changes in cell morphology and release of tumor necrosis factor-alpha (Levy et al., *Annu. Rev. Immunol.*, 1998, 16, 89-109).

CD81 has been identified as a potential receptor for entry of the hepatitis C virus via the HCV E2 envelope protein, a finding which may contribute to the elucidation of the pathogenesis of HCV-associated diseases and aid in the development of new therapeutic strategies directed at virus binding interference (Pileri et al., *Science*, 1998, 282, 938-941). Studies by Bisceglie support the CD81-HCV E2 interaction as a target for future antiviral therapies (Bisceglie, *Am. J. Med.*, 1999, 107, 45S-48S). Disclosed and claimed in PCT publication WO 01/58459 is a substance inhibiting the binding of HCV E2/NS1 protein to cells having the potential to be infected with HCV, including cells expressing CD81 (Shibui et al., 2001).

RTSP-0443

-3-

Inhibition of protein-protein interactions involving CD81 interactions are well known in the art. For example, Fleming et al. have shown that antibodies recognizing CD81 inhibit mast cell degranulation mediated by the high affinity receptor of immunoglobulin E, indicating an unsuspected calcium-independent pathway of antigen receptor regulation which is accessible to engagement by membrane proteins and on which novel therapeutic approaches to allergic diseases could be based (Fleming et al., *J. Exp. Med.*, **1997**, 186, 1307-1314).

CD81-null mice undergo normal development of thymocytes and express normal quantities of T-cells and B-cells but they express lower levels of CD19, indicating that CD81 is not required for maturation of T-cells but is important for optimal expression of CD19 on B-cells and optimal stimulation of antibody production (Maecker and Levy, *J. Exp. Med.*, **1997**, 185, 1505-1510). CD81-deficient mice were reported to have diminished allergen-induced airway hyper-reactivity due to alterations in local cytokine production, indicating that further dissection of the immunological and physiological defect of these mice may provide targets for the development of new anti-asthma therapies (Deng et al., *J. Immunol.*, **2000**, 165, 5054-5061).

CD81-deficient mice were also reported to have an altered sensitivity to the neurobehavioral effects of cocaine, indicating that CD81 is involved in mediating the acute and/or long-term neural responsiveness to cocaine (Michna et al., *Mol. Brain Res.*, **2001**, 90, 68-74). Furthermore, upregulation of CD81 expression in the nucleus accumbens of cocaine-treated rats provides support for the latter investigation (Brenz Verca et al., *Mol. Cell. Neurosci.*, **2001**, 17, 303-316).

Since CD81 acts as an HCV receptor and immune regulator and is known to influence responsiveness to cocaine, its selective inhibition may be a potential strategy with which to derive treatments for viral infections, autoimmune disorders and cocaine addiction.

Disclosed and claimed in PCT publication WO 98/25647 is a method of inhibiting passive cutaneous anaphylaxis in a mammal comprising administering to the mammal an effective amount of an



RTSP-0443

-4-

agent which inhibits CD81-mediated signal transduction (Fleming and Kinet, 1998).

Currently, there are no known therapeutic agents that effectively inhibit the synthesis of CD81. To date, investigative strategies aimed at modulating CD81 expression have involved the use of antibodies and gene knock-outs in mice. Consequently, there remains a long felt need for agents capable of effectively inhibiting CD81 function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of expression of CD81.

The present invention provides compositions and methods for modulating expression of CD81.

#### **SUMMARY OF THE INVENTION**

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding CD81, and which modulate the expression of CD81. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of CD81 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of CD81 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

#### **DETAILED DESCRIPTION OF THE INVENTION**

The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding CD81, ultimately

RTSP-0443

-5-

modulating the amount of CD81 produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding CD81. As used herein, the terms "target nucleic acid" and "nucleic acid encoding CD81" encompass DNA encoding CD81, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of CD81. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding CD81. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a

RTSP-0443

-6-

preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding CD81, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also

RTSP-0443

-7-

a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

It is also known in the art that alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and extronic regions.

RTSP-0443

-8-

Upon excision of one or more exon or intron regions or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

It is also known in the art that variants can be produced through the use of alternative signals to start or stop transcription and that pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each

RTSP-0443

-9-

other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other

RTSP-0443

-10-

antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

Expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, **2000**, 480, 17-24; Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, **2000**, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, **1999**, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U. S. A.*, **2000**, 97, 1976-81), protein arrays and proteomics (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16; Jungblut, et al., *Electrophoresis*, **1999**, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16; Larsson, et al., *J. Biotechnol.*, **2000**, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., *Anal. Biochem.*, **2000**, 286, 91-98; Larson, et al., *Cytometry*, **2000**, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, **2000**, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, **1998**, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, **1999**, 35, 1895-904) and mass spectrometry methods (reviewed in (To, *Comb. Chem. High Throughput Screen*, **2000**, 3, 235-41).

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses.

RTSP-0443

-11-

Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines.



RTSP-0443

-12-

Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside

RTSP-0443

-13-

residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenhydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

RTSP-0443

-14-

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylinino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>m</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl,

RTSP-0443

-15-

substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, **1995**, *78*, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, also described in examples hereinbelow.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH<sub>2</sub>-)<sub>n</sub> group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137;

RTSP-0443

-16-

5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ( $-C\equiv C-CH_3$ ) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United

RTSP-0443

-17-

States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention

RTSP-0443

-18-

include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, *86*, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1994**, *4*, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, *660*, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1993**, *3*, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, *20*, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, **1991**, *10*, 1111-1118; Kabanov et al., *FEBS Lett.*, **1990**, *259*, 327-330; Svinarchuk et al., *Biochimie*, **1993**, *75*, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, **1995**, *36*, 3651-3654; Shea et al., *Nucl. Acids Res.*, **1990**, *18*, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, **1995**, *14*, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, **1995**, *36*, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, **1995**, *1264*, 229-237), or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Croke et al., *J. Pharmacol. Exp. Ther.*,

RTSP-0443

-19-

1996, 277, 923-937. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an



RTSP-0443

-20-

oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the

RTSP-0443

-21-

phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

RTSP-0443

-22-

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid,

RTSP-0443

-23-

glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder

RTSP-0443

-24-

which can be treated by modulating the expression of CD81 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding CD81, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding CD81 can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of CD81 in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments,

RTSP-0443

-25-

lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof,

RTSP-0443

-26-

bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate,. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail

RTSP-0443

-27-

in United States applications 08/886,829 (filed July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams.



RTSP-0443

-28-

Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are

RTSP-0443

-29-

comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in

RTSP-0443

-30-

*Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may

RTSP-0443

-31-

readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to

RTSP-0443

-32-

form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, *in: Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, *in Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and

RTSP-0443

-33-

consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of

RTSP-0443

-34-

oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

RTSP-0443

-35-

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured,



RTSP-0443

-36-

releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, **1992**, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, **1992**, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, **1992**, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and

RTSP-0443

-37-

cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G<sub>M1</sub>, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G<sub>M1</sub> or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-

RTSP-0443

-38-

dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, **1980**, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, **1984**, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, **1990**, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, **1990**, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in

RTSP-0443

-39-

liposomes. U.S. Patent No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters,

RTSP-0443

-40-

polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a

RTSP-0443

-41-

penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

RTSP-0443

-42-

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium

RTSP-0443

-43-

ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic



RTSP-0443

-44-

acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyanostilbene-2,2'-disulfonic acid (Miyao *et al.*, *Antisense Res. Dev.*, **1995**, 5, 115-121; Takakura *et al.*, *Antisense & Nucl. Acid Drug Dev.*, **1996**, 6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch,

RTSP-0443

-45-

sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of

RTSP-0443

-46-

the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide).

RTSP-0443

-47-

Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on  $EC_{50}$ s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the

RTSP-0443

-48-

oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

RTSP-0443

-49-

**EXAMPLES****Example 1****Nucleoside Phosphoramidites for Oligonucleotide Synthesis  
Deoxy and 2'-alkoxy amidites**

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, **1993**, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

**2'-Fluoro amidites****2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, **1993**, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

RTSP-0443

-50-

**2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

**2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

**2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'-phosphoramidites.

**2'-O-(2-Methoxyethyl) modified amidites**

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

**2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenyl-carbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas

RTSP-0443

-51-

to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

#### **2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

#### **2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of



RTSP-0443

-52-

dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with  $\text{CH}_3\text{CN}$  (200 mL). The residue was dissolved in  $\text{CHCl}_3$  (1.5 L) and extracted with 2x500 mL of saturated  $\text{NaHCO}_3$  and 2x500 mL of saturated  $\text{NaCl}$ . The organic phase was dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with  $\text{EtOAc}$ /hexane/acetone (5:5:1) containing 0.5%  $\text{Et}_3\text{NH}$ . The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in  $\text{CHCl}_3$  (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated  $\text{NaCl}$ . The water layers were back extracted with 200 mL of  $\text{CHCl}_3$ . The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using  $\text{EtOAc}$ /hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

RTSP-0443

-53-

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in  $\text{CH}_3\text{CN}$  (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in  $\text{CH}_3\text{CN}$  (1 L), cooled to  $-5^\circ\text{C}$  and stirred for 0.5 h using an overhead stirrer.  $\text{POCl}_3$  was added dropwise, over a 30 minute period, to the stirred solution maintained at  $0-10^\circ\text{C}$ , and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of  $\text{NaHCO}_3$  and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and  $\text{NH}_4\text{OH}$  (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with  $\text{NH}_3$  gas was added and the vessel heated to  $100^\circ\text{C}$  for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

RTSP-0443

-54-

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra(isopropyl)-phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites****2'-(Dimethylaminooxyethoxy) nucleoside amidites**

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs.

RTSP-0443

-55-

Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine**  
O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained

RTSP-0443

-56-

for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

RTSP-0443

-57-

**5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine**

2'-O-([2-phthalimidooxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry  $\text{CH}_2\text{Cl}_2$  (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at  $-10^\circ\text{C}$ . to  $0^\circ\text{C}$ . After 1 h the mixture was filtered, the filtrate was washed with ice cold  $\text{CH}_2\text{Cl}_2$ , and the combined organic phase was washed with water, brine and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

**5'-O-tert-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine**

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at  $10^\circ\text{C}$  under inert atmosphere. The reaction mixture was stirred for 10 minutes at  $10^\circ\text{C}$ . After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ). Aqueous  $\text{NaHCO}_3$  solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to  $10^\circ\text{C}$  in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at  $10^\circ\text{C}$  for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5%  $\text{NaHCO}_3$ ,

RTSP-0443

-58-

(25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in  $\text{CH}_2\text{Cl}_2$  to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in  $\text{CH}_2\text{Cl}_2$  to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

**5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine**

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over  $\text{P}_2\text{O}_5$  under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in  $\text{CH}_2\text{Cl}_2$  (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

RTSP-0443

-59-

**5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N',N'-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P.



RTSP-0443

-60-

D., Guinasso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-([2-phthalamidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

RTSP-0443

-61-

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with  $\text{CH}_2\text{Cl}_2$  (2x200 mL). The combined  $\text{CH}_2\text{Cl}_2$  layers are washed with saturated  $\text{NaHCO}_3$  solution, followed by saturated  $\text{NaCl}$  solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using  $\text{MeOH}:\text{CH}_2\text{Cl}_2:\text{Et}_3\text{N}$  (20:1, v/v, with 1% triethylamine) gives the title compound.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyluridine (2.17 g, 3 mmol) dissolved in  $\text{CH}_2\text{Cl}_2$  (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

**Example 2****Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from

RTSP-0443

-62-

the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance,

RTSP-0443

-63-

alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### Example 4

##### PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

#### Example 5

##### Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

##### [2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated

RTSP-0443

-64-

DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)]**

**Chimeric Phosphorothioate Oligonucleotides**

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

**[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides**

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the

RTSP-0443

-65-

chimeric structures and sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

#### Example 6

##### Oligonucleotide Isolation

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* **1991**, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

#### Example 7

##### Oligonucleotide Synthesis - 96 Well Plate Format

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA,

RTSP-0443

-66-

or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature ( $55-60^\circ\text{C}$ ) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

#### **Example 8**

##### **Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

#### **Example 9**

##### **Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 4 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

RTSP-0443

-67-

## T-24 cells:

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

## A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

## NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

## HEK cells:



RTSP-0443

-68-

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If

RTSP-0443

-69-

60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

#### Example 10

##### Analysis of oligonucleotide inhibition of CD81 expression

Antisense modulation of CD81 expression can be assayed in a variety of ways known in the art. For example, CD81 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

Protein levels of CD81 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to CD81 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can

RTSP-0443

-70-

be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

**Example 11****Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

RTSP-0443

-71-

**Example 12****Total RNA Isolation**

Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 µL cold PBS. 100 µL Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100 µL of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 µL water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 µL water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

RTSP-0443

-72-

**Example 13****Real-time Quantitative PCR Analysis of CD81 mRNA Levels**

Quantitation of CD81 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

RTSP-0443

-73-

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN<sup>TM</sup> buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD<sup>TM</sup>, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD<sup>TM</sup>, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen<sup>TM</sup> (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen<sup>TM</sup> RNA quantification reagent from

RTSP-0443

-74-

Molecular Probes. Methods of RNA quantification by RiboGreen™ are taught in Jones, L.J., et al, *Analytical Biochemistry*, 1998, 265, 368-374.

In this assay, 175 µL of RiboGreen™ working reagent (RiboGreen™ reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25µL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

Probes and primers to human CD81 were designed to hybridize to a human CD81 sequence, using published sequence information (GenBank accession number M33680.1, incorporated herein as SEQ ID NO:3). For human CD81 the PCR primers were:

forward primer: CAGATCGCCAAGGATGTGAA (SEQ ID NO: 4)

reverse primer: GCGTTGTTGGCGTCATCA (SEQ ID NO: 5) and the PCR

probe was: FAM-AGTTCTATGACCAGGCCCTACAGCAGGC-TAMRA

(SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO:7)

reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO:8) and the PCR

probe was: 5' JOE-CAAGCTTCCCCTTCTCAGCC-TAMRA 3' (SEQ ID NO: 9)

where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

#### Example 14

##### Northern blot analysis of CD81 mRNA levels

Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham

RTSP-0443

-75-

Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions.

To detect human CD81, a human CD81 specific probe was prepared by PCR using the forward primer CAGATCGCCAAGGATGTGAA (SEQ ID NO: 4) and the reverse primer GCGTTGTTGGCGTCATCA (SEQ ID NO: 5). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### Example 15

##### **Antisense inhibition of human CD81 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human CD81 RNA, using published sequences (GenBank accession number M33680.1, incorporated herein as SEQ ID NO: 3; GenBank accession number AA595584.1 representing a 3' extension from SEQ ID NO: 3, the complement of which is incorporated herein as SEQ ID NO: 10; and residues 59000-880500 of GenBank accession NT\_009368.4 representing a partial genomic sequence of CD81, the complement of which is incorporated herein as SEQ ID NO: 11). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds.



RTSP-0443

-76-

All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human CD81 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

Inhibition of human CD81 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
193983	5'UTR	3	1	ggcgcctttccagcaccaatgg	0	13
193990	Coding	3	291	agccagccagagaagacgaaat	37	14
194022	3'UTR	3	1221	cctggctgagcagggcagagc	39	15
199577	Coding	3	246	gcacttggtgcagccctcca	76	16
199578	Coding	3	272	ttgaagacgaagagcaggta	40	17
199579	Coding	3	277	cgaatttgaagacgaagagc	0	18
199580	Coding	3	286	gccagaagacgaatttgaag	26	19
199581	Coding	3	296	cctccagccagccagaagac	43	20
199582	Coding	3	304	ggatcacgcctccagccagc	81	21
199583	Coding	3	320	cacagggccacacccaggat	62	22
199584	Coding	3	375	cttgtctcccagctccagat	41	23
199585	Coding	3	402	gcctacatagaagggtgttgg	48	24
199586	Coding	3	410	atgtagatgcctacatagaa	22	25
199587	Coding	3	471	gatggccccgtagcagccca	76	26
199588	Coding	3	497	gtccccagcaggcactggga	62	27
199589	Coding	3	536	acctcacaggcaaacaggat	26	28
199590	Coding	3	564	gttgacaaagccccagatgc	78	29
199591	Coding	3	587	acatccttggcgatctggtc	69	30
199592	Coding	3	592	gcttcacatccttggcgatc	44	31
199593	Coding	3	597	gaactgcttcacatccttgg	81	32
199594	Coding	3	602	tcatagaactgcttcacatc	3	33
199595	Coding	3	607	cctggctcatagaactgcttc	57	34
199596	Coding	3	665	gtcttcaccacagccttggc	63	35
199597	Coding	3	678	cgtctcgtggaaggtcttca	81	36
199598	Coding	3	747	gggacacaaattgttcttga	30	37
199599	Coding	3	753	gcccggaggacacaaattgt	69	38
199600	Coding	3	781	ccttgaagaggttgctgatg	47	39
199601	Coding	3	790	ggcagtcctccttgaagagg	55	40

RTSP-0443

-77-

199602	Coding	3	830	aggtacagcttccccggagaa	65	41
199603	Coding	3	836	ccgatgaggtacagcttccc	75	42
199604	Coding	3	841	caatgccgatgaggtacagc	68	43
199605	Coding	3	848	atggcagcaatgccgatgag	38	44
199606	Coding	3	857	gcgaccacgatggcagcaat	53	45
199607	Coding	3	864	gatcacagcgaccacgatgg	52	46
199608	Coding	3	884	aggatcatctcgaagatcat	25	47
199609	Coding	3	896	aggaccatgctcaggatcat	78	48
199610	Coding	3	902	cagcacagcaccatgctcag	41	49
199611	Coding	3	907	tgccacagcacagcaccatg	0	50
199612	Coding	3	912	ccggatgccacagcacagca	79	51
199613	Coding	3	917	ctgttccggatgccacagca	71	52
199614	Coding	3	922	cggagctgttccggatgcca	77	53
199615	Coding	3	927	gtacacggagctgttccgga	70	54
199616	Stop Codon	3	935	gggcctcagtacacggagct	55	55
199617	3'UTR	3	962	gcagaggtccctgtggccag	59	56
199618	3'UTR	3	1015	tacacaggcggatgagccc	69	57
199619	3'UTR	3	1048	tacgtgtagcagagtaatac	56	58
199620	3'UTR	3	1060	aagtaaaaaggctacgtgta	73	59
199621	3'UTR	3	1098	aggtaacaggaaagttcaga	74	60
199622	3'UTR	3	1112	catcagccctgaaaaggtaa	46	61
199623	3'UTR	3	1136	ctcatcacagccacctacat	2	62
199624	3'UTR	3	1184	gcagaaggaccctgcccctc	50	63
199625	3'UTR	3	1205	gagcaccctgggaccccagg	74	64
199626	3'UTR	3	1210	aggcagagcaccctgggacc	68	65
199627	3'UTR	3	1242	gcgagtggctcccaggagag	67	66
199628	3'UTR	3	1261	ggccaagctgagttctctggg	34	67
199629	3'UTR	3	1309	gtgcagcccacaggacgggc	76	68
199630	3'UTR	3	1327	gaggggaacaagggtgagctgt	10	69
199631	3'UTR	3	1355	cacagactcggctctcgaac	2	70
199632	3'UTR	3	1369	aggcagagagtgccacaga	71	71
199633	3'UTR	3	1375	gcataaggcagagagtgcc	68	72
199634	3'UTR	3	1393	gtgttagaaaaggacaggtgc	19	73
199635	3'UTR	3	1409	tacagttgaaggcgacgtgt	59	74
199636	3'UTR	3	1415	tgtgattacagttgaaggcg	65	75
199637	3'UTR	3	1430	acggagtcaggatgtttgtga	94	76
199638	3'UTR	3	1440	ttattaaatgacggagtcag	29	77
199639	3'UTR	3	1447	tccttctttattaaatgacg	80	78
199640	Intron 1	11	9288	agcgcaaccagtggaactggc	61	79
199641	Intron 1	11	10090	ccaggcctctgccgaggggtt	40	80
199648	Intron 2	11	15557	ggcctcagcacagcctgtga	71	81
199649	Intron 3	11	17485	acctccccagaaattctagg	31	82
199650	Intron : Exon Junction	11	18159	aggtgaagaactggagccgg	36	83
199651	Exon: Intron Junction	11	18234	ccaggctcacctggctccttg	0	84
199652	Intron: Exon Junction	11	19056	agcagtcgaagctgcagagag	0	85
199653	Exon: Intron Junction	11	19902	gcccgcctcaccatgatcaca	31	86
199654	Exon 1	11	657	ctggctgggtggccggcaaag	18	87
199655	Exon 8	11	20210	gtaacaggaaagttcagaac	73	88
199656	Exon 8	11	20224	tcagccctgaaaaggtaaca	50	89
199657	3'UTR	10	577	tttttgactgcacaggcctg	2	90

RTSP-0443

-78-

As shown in Table 1, SEQ ID NOS 14, 15, 16, 17, 20, 21, 22, 23, 24, 26, 27, 29, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 48, 49, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 63, 64, 65, 66, 67, 68, 71, 72, 74, 75, 76, 78, 79, 80, 81, 82, 83, 86, 88 and 89 demonstrated at least 30% inhibition of human CD81 expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

**Example 16****Western blot analysis of CD81 protein levels**

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to CD81 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

**What is claimed is:**

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding CD81, wherein said compound specifically hybridizes with said nucleic acid molecule encoding CD81 and inhibits the expression of CD81.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 14, 15, 16, 17, 20, 21, 22, 23, 24, 26, 27, 29, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 48, 49, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 63, 64, 65, 66, 67, 68, 71, 72, 74, 75, 76, 78, 79, 80, 81, 82, 83, 86, 88 or 89.

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding CD81.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

RTSP-0443

-80-

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of CD81 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of CD81 is inhibited.

16. A method of treating an animal having a disease or condition associated with CD81 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of CD81 is inhibited.

17. The method of claim 16 wherein the disease or condition is an inflammatory disorder.

18. The method of claim 16 wherein the disease or condition is an infection.

19. The method of claim 18 wherein the infection is selected from the group consisting of viral, bacterial and parasitic.

20. The method of claim 16 wherein the disease or condition is characterized by chemical dependency.

## SEQUENCE LISTING

<110> Mark J. Graham  
Kenneth Dobie  
Isis Pharmaceuticals, Inc.

<120> ANTISENSE MODULATION OF CD81 EXPRESSION

<130> RTSP-0443

<150> 10/006,430  
<151> 2001-12-10

<160> 90

<210> 1  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1  
tccgtcatcg ctcctcaggg 20

<210> 2  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2  
atgcattctg cccccaagga 20

<210> 3  
<211> 1496  
<212> DNA  
<213> Homo sapiens

<220>

<221> CDS  
<222> (239)...(949)

<400> 3  
ccattgtgct ggaaaggcgc gcaacggcgg cgacggcggc gaccccaccg cgcattcctgc 60  
caggcctccg cgcccagccg cccacgcgcc cccgcgcccc gcgccccgac cttttcttcg 120  
cgcccccgcc cctcgggccc ccaggccccc ttgccggcca cccgccaggc cccgcgccgg 180

cccgccccgcc gccaggacc ggcccgcgcc ccgcaggccg cccgcccgcc gcgcccgc 238  
  
 atg gga gtg gag ggc tgc acc aag tgc atc aag tac ctg ctc ttc gtc 286  
 Met Gly Val Glu Gly Cys Thr Lys Cys Ile Lys Tyr Leu Leu Phe Val  
 1 5 10 15  
  
 ttc aat ttc gtc ttc tgg ctg gct gga ggc gtg atc ctg ggt gtg gcc 334  
 Phe Asn Phe Val Phe Trp Leu Ala Gly Gly Val Ile Leu Gly Val Ala  
 20 25 30  
  
 ctg tgg ctc cgc cat gac ccg cag acc acc aac ctc ctg tat ctg gag 382  
 Leu Trp Leu Arg His Asp Pro Gln Thr Thr Asn Leu Leu Tyr Leu Glu  
 35 40 45  
  
 ctg gga gac aag ccc gcg ccc aac acc ttc tat gta ggc atc tac atc 430  
 Leu Gly Asp Lys Pro Ala Pro Asn Thr Phe Tyr Val Gly Ile Tyr Ile  
 50 55 60  
  
 ctc atc gct gtg ggc gct gtc atg atg ttc gtt ggc ttc ctg ggc tgc 478  
 Leu Ile Ala Val Gly Ala Val Met Met Phe Val Gly Phe Leu Gly Cys  
 65 70 75 80  
  
 tac ggg gcc atc cag gaa tcc cag tgc ctg ctg ggg acg ttc ttc acc 526  
 Tyr Gly Ala Ile Gln Glu Ser Gln Cys Leu Leu Gly Thr Phe Phe Thr  
 85 90 95  
  
 tgc ctg gtc atc ctg ttt gcc tgt gag gtg gcc gcc ggc atc tgg ggc 574  
 Cys Leu Val Ile Leu Phe Ala Cys Glu Val Ala Ala Gly Ile Trp Gly  
 100 105 110  
  
 ttt gtc aac aag gac cag atc gcc aag gat gtg aag cag ttc tat gac 622  
 Phe Val Asn Lys Asp Gln Ile Ala Lys Asp Val Lys Gln Phe Tyr Asp  
 115 120 125  
  
 cag gcc cta cag cag gcc gtg gtg gat gat gac gcc aac aac gcc aag 670  
 Gln Ala Leu Gln Gln Ala Val Val Asp Asp Asp Ala Asn Asn Ala Lys  
 130 135 140  
  
 gct gtg gtg aag acc ttc cac gag acg ctt gac tgc tgt ggc tcc agc 718  
 Ala Val Val Lys Thr Phe His Glu Thr Leu Asp Cys Cys Gly Ser Ser  
 145 150 155 160  
  
 aca ctg act gct ttg acc acc tca gtg ctc aag aac aat ttg tgt ccc 766  
 Thr Leu Thr Ala Leu Thr Thr Ser Val Leu Lys Asn Asn Leu Cys Pro  
 165 170 175  
  
 tcg ggc agc aac atc atc agc aac ctc ttc aag gag gac tgc cac cag 814  
 Ser Gly Ser Asn Ile Ile Ser Asn Leu Phe Lys Glu Asp Cys His Gln  
 180 185 190  
  
 aag atc gat gac ctc ttc tcc ggg aag ctg tac ctc atc ggc att gct 862  
 Lys Ile Asp Asp Leu Phe Ser Gly Lys Leu Tyr Leu Ile Gly Ile Ala  
 195 200 205  
  
 gcc atc gtg gtc gct gtg atc atg atc ttc gag atg atc ctg agc atg 910  
 Ala Ile Val Val Ala Val Ile Met Ile Phe Glu Met Ile Leu Ser Met  
 210 215 220

gtg ctg tgc tgt ggc atc cgg aac agc tcc gtg tac tga ggccccgcag 959  
 Val Leu Cys Cys Gly Ile Arg Asn Ser Ser Val Tyr  
 225 230 235

ctctggccac agggacctct gcagtgcccc ctaagtgacc cggacacttc cgagggggcc 1019  
 atcaccgcct gtgtatataa cgtttccggt attactotgc tacacgtagc ctttttaactt 1079  
 ttgggggtttt gtttttggtc tgaactttcc tgttaccttt tcagggctga tgtcacatgt 1139  
 aggtggcgtg tatgagtgga gacgggcctg ggtcttgggg actggagggc aggggtcctt 1199  
 ctgcccctgg ggtcccaggg tgctctgcct gctcagccag gcctctcctg ggagccactc 1259  
 gccagagac tcagcttggc caacttgggg ggctgtgtcc acccagcccg cccgtcctgt 1319  
 gggctgcaca gctcaccttg ttccctcctg ccccggttcg agagccgagt ctgtgggcac 1379  
 tctctgcctt catgcacctg tcctttctaa cacgtcgctt tcaactgtaa tcacaacatc 1439  
 ctgactcogt catttaataa agaaggaaca tcaggcatgc taaaaaaaaa aaaaaaa 1496

<210> 4  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 4  
 cagatcgcca aggatgtgaa 20

<210> 5  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5  
 gcgttggttg cgtcacatca 18

<210> 6  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Probe



<400> 6  
 agttctatga ccaggcccta cagcaggc 28

<210> 7  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 7  
 gaaggtgaag gtcggagtc 19

<210> 8  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 8  
 gaagatggtg atgggatttc 20

<210> 9  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Probe

<400> 9  
 caagcttccc gttctcagcc 20

<210> 10  
 <211> 626  
 <212> DNA  
 <213> Homo sapiens

<220>

<400> 10  
 atccgagcct gtgcttgctg ggcacgga cagctcgtgt attgagtccc gcagctttgg 60  
 ccacaagaac ttctgcagtg cccctaagt gaccgga cttccgagg gaccatcacc 120  
 gcctgtgtat ataacgtttc cggattact ctgctacacg tagcctttt acttttggg 180  
 ttttgttttt gttccgaact ttctgttac cttttcagg ctgacgtcac atgtaggtgg 240

```

cgtgtatgag tggagacggg cctgggtctt ggggactgga gggcaggggt ccttctgccc 300
tggggtccca ggggtgctctg cctgctcagc caggcctctc ctgggagcca ctgcccaga 360
gactcagctt ggccaacttg gggggctgtg tccaccagc ccgccgtcc tgtgggctgc 420
acagctcacc ttgttccctc ctgccccggt tcgagagccg agtctgtggg cactctctgc 480
cttcatgcac ctgtcctttc taacacgtcg cttcaactg taatcacaac atcctgactc 540
cgtcatttaa taaagaagga acatcaggca tgctaccag cctgtgcagt caaaaaaaaa 600
aaaaaaaaa aaaaaaaaaa aaaaaa 626

```

<210> 11

<211> 21501

<212> DNA

<213> Homo sapiens

<220>

<221> exon

<222> (518)...(812)

<223> exon 1

<221> exon:intron junction

<222> (812)...(813)

<223> exon 1:intron 1

<221> intron

<222> (813)...(13608)

<223> intron 1

<221> intron:exon junction

<222> (13608)...(13609)

<223> intron 1:exon 2

<221> exon

<222> (13609)...(13723)

<223> exon 2

<221> exon:intron junction

<222> (13723)...(13724)

<223> exon 2:intron 2

<221> intron

<222> (13724)...(17291)

<223> intron 2

<221> intron:exon junction

<222> (17291)...(17292)

<223> intron 2:exon 3

<221> exon

<222> (17292)...(17389)

<223> exon 3

<221> exon:intron junction

<222> (17389)...(17390)  
<223> exon 3:intron 3

<221> intron  
<222> (17390)...(18168)  
<223> intron 3

<221> intron:exon junction  
<222> (18168)...(18169)  
<223> intron 3:exon 4

<221> exon  
<222> (18169)...(18243)  
<223> exon 4

<221> exon:intron junction  
<222> (18243)...(18244)  
<223> exon 4:intron 4

<221> intron  
<222> (18244)...(18612)  
<223> intron 4

<221> intron:exon junction  
<222> (18612)...(18613)  
<223> intron 4:exon 5

<221> exon  
<222> (18613)...(18717)  
<223> exon 5

<221> exon:intron junction  
<222> (18717)...(18718)  
<223> exon 5:intron 5

<221> intron  
<222> (18718)...(19065)  
<223> intron 5

<221> intron:exon junction  
<222> (19065)...(19066)  
<223> intron 5:exon 6

<221> exon  
<222> (19066)...(19167)  
<223> exon 6

<221> exon:intron junction  
<222> (19167)...(19168)  
<223> exon 6:intron 6

<221> intron  
<222> (19168)...(19824)  
<223> intron 6

<221> intron:exon junction  
<222> (19824)...(19825)  
<223> intron 6:exon 7

<221> exon  
 <222> (19825)...(19911)  
 <223> exon 7

<221> exon:intron junction  
 <222> (19911)...(19912)  
 <223> exon 7:intron 7

<221> intron  
 <222> (19912)...(20000)  
 <223> intron 7

<221> intron:exon junction  
 <222> (20000)...(20001)  
 <223> intron 7:exon 8

<221> exon  
 <222> (20001)...(20610)  
 <223> exon 8

<400> 11  
 catgcctgcc caggttccag ccccgagag caatgtgagc aaagcttgct gtctttgcaa 60  
 agccaaccac tgtggcatca actccttcag gaagccctcc cggattgtcc aaggtgctca 120  
 cctccttttg ggagccctcc cagattgtcc aaggtgcttg agggagggag gaatgggttg 180  
 ttctcccggc accggggctg cactcctggg cagacgtgc atgcctgtcc tcaggcgcgg 240  
 ccctgctgcc acccccttg gggtcggag cgcgacagca gcttg'gggac gcctcccgcg 300  
 cccagcacgg tgcacctggg ccctgaggtc ctggccgaaa cgcgccaagt tgggggtagg 360  
 tgcagcgacc ccatacccct cggtgcgcg ccctggcggc aggagggcggg gccgggggcg 420  
 gggcgtgagc tggccggggg cggggcctat ggaggggcgg gaccgcggcg ccctataagt 480  
 actgcggagc gcaggcgcgc gcccgccag agagcgagcg cgcaacggcg gcgacggcgg 540  
 cgaccccacc gcacatcctg ccaggcctcc ggcgcccagg gcgcacggcg cgccccctg 600  
 ccggcgggcc ctgcgccc atcttggcgc ccccgcccgg tcggcccgc agggcccttt 660  
 gccggccacc agccaggccc cgcgcggcc cgcccgcgc ccaggaccgg ccgcgcgcc 720  
 gcaggccgc cgccgccgc gccgccatgg gagtggaggg ctgcaccaag tgcataagt 780  
 acctgctctt cgtcttcaat ttcgtcttct gggtaagggc tgcgcggggg gccggggcgg 840  
 gagggggcag gcacacactc cacgttgggc aggtcccgc gcagcgtgct agggcccgcg 900  
 ggcgcagcgc gggccgcgaa gttgtggggc cacctgtggg ctccaggagc ggggtggggg 960  
 gtcgcccggg gccaccgcgc ccccgacat tggggctgag ggctgcgagc cgagtttcgg 1020  
 ggcctctgtg ctggggggc cacctctgc gccgggccgg ggcttctggg ggccgcgggg 1080  
 cagttccgc tgtggtggtg atgggtgcgg tggtcgcggg tcgggaccgc agtaccgcgc 1140

cgccccctcag ctaaggaggg gcoctgcgcgg gtccctggcc gcggattccg gactgctgct 1200  
 tcgcgggggac gagggggggg ctgcggggcg ggactcctgg cgccccgccc ccatgagctc 1260  
 atcaagagcc gccgcccctg gatggtgggg cgggggcgca cactttgccc gaggttgggg 1320  
 gcgatccgcc tcaactcttc cccagcccag ctcaactctcc aatctgcggt caccaccoga 1380  
 gaccttcctg ggggtcgcgc ctaaaaggag cgcagactcc cgccgggatg gcccagaagc 1440  
 tggggtgcgc gcaccctggc cgtccctgcc tgggagccga tctccctctc ctcaccaga 1500  
 cacgttcag cgagggcctc ctcccagaag ggctctggag gcctcgcagg agtggggatc 1560  
 ccgcggttct gagttggcac aaggaagaga gtggcaccag gggcctggag tggatggcag 1620  
 ggtccgggag tggggccgct gctttgcaag aggggcccc acgctgggca tctttgggtg 1680  
 ccagcgtggg tggaggaggg tcttttctg agaatggctt tctcctgacc gcagtctttg 1740  
 ctgctgggaa gtgactgatg ggctttcgcc ttttgtttcc atttcctgtc ggtgttagaa 1800  
 ttggggaggg ggtggaaatc ctttcttggc ctggaaggac tggagtgggt gtccatggcc 1860  
 gcggcctccc cgtggccacg ccctgggca tagactgcaa gcccctcccc gtgccccca 1920  
 ggctgtcacc cccttctcgt ggaagactcg gctgatgtcc cagtggaccg agtgtttctc 1980  
 aagttgaggc agggagggca aactttttaa atggccctg gagccagtgt gtgggaccag 2040  
 agacatctgt ttcccatctg gacggctgag gatcccagtg cggatgatta tttggagggg 2100  
 gaaggacgga ggctgaactg aactctcagc tgggagatga gtggggcagt cacatccac 2160  
 cttccccaag cggggctgtt ctgcacagcc tgcttgggac gctggtggga gtcactgtgg 2220  
 ctttcggcac tgccctggca gtgggggag ctaggccatt tgggaggggc tcgctttccc 2280  
 caggccgggc cctgggacct cagccgttgc ttagtggtgg cctgcttcag cccaggcatg 2340  
 tgggagaggc accagacaca ggatgtccct ctgccagccc ctgaagcccc gtccctgac 2400  
 gaggcgagtg tggacctggg ggtgggggct gagggagact gtggacctgg gggtgggggc 2460  
 tgaaggaagg tgtggacctg ggggcagggg ccgagggaag gtgtaggcct ggggtagta 2520  
 ggggctgagg gagagtgtg acctgggagt aggggctgag ggaggggtga ggcctggggg 2580  
 tgggggctga gggagagtgt ggacctggg gtaggggctg agggagagtg tggacctggg 2640  
 ggtgggggtt gagggagggt gtggacctgg gggcaggggc tgaggagag tgtggacct 2700  
 ggggcagagg ctgaagggga gtcacgggag gggacttctc cggaggtgga tttttgctct 2760  
 ctggacggtg tgtcagcact gggtagccc ctctgcctg cccaggctga gaggtctccc 2820  
 tggcagcccc ctgggagtgt cgccagggcg ggcctggaag tttcccaggc agctgggggtg 2880  
 gagacctgac acatcccaag ggtgcttgtt attaaggctc aaggaaatgt ctctgaggcc 2940

tcaccgctcc tctccccagg gcttgctccc tgcaaagcat tgagaactga gtccgtccac 3000  
 agtcactgtg gaccacacca tccactgggg ctgagtggtg gccagcaatg ccaggctggg 3060  
 tgagggtggg ttggtgggca ccacctgggt ggacccccct ccacctgggt gtcgcagggt 3120  
 gtgtggctga gagcacagt ccatgggctt gggcctcctt ggtggagtcc ccaacacact 3180  
 gctctggtcc tgggcctcgg ccttccccgt ctgagtggtg ggcccacagt gagcctacct 3240  
 cctggtggtg ttggtggatt tgctgacatg cctgagtgtt gacagggggc ttggtgcagg 3300  
 aagggctcag ggctgggtg ttggccaggg gtccaaaggg acctctgcct cagagagccc 3360  
 agcccagaca ggcaggatgt gcagtgggga aggggctgcg ggaaccctgc aggggtccaga 3420  
 aggacacagt gcagtcctgt gggctctggg gaggtggtg gggaggaggt tgacaatgga 3480  
 tatctgggtg gggcacttgt tagaagttcc attttagaga ggaaagaggc cttgcctgtg 3540  
 ggagaaggca gctggggtag cctgacctct tcccaggaa ggagcccaca cacacacgca 3600  
 caggcactca cacacacgaa tgtgcacaca cacacactcc caccttcaca cacttcaca 3660  
 ctcttgctgt ctccttccc aagccaaggt gcgaggggga aggtctgggc agcatgcacc 3720  
 tgccgctga ccgctttggg ggccagtga aactgggctc cctgggtgcg cggcgggccc 3780  
 aagcaggag gacattgcag atgccctggc caagcagcgt ggaaatcctg tcccttgggt 3840  
 ggtctcga gcctccatca gagggcgctg gcacctgaga cccacctgct gccaggagca 3900  
 gggcaggaga gtttgtgtcc cgggacaggg aactggcctg tgggagcctt gccttccca 3960  
 tctgtgtaat ggatataaga gtcttctcct cgggggctgg ccagggagtc cagaagaggt 4020  
 gtcaccagtc ccgcaggga gaagagcggg gtccccgcc tgggactggc tgcctcccca 4080  
 agctaattgca gctggtagcc acctcccagt ggcagggcag ccaaaccg cgggaaaga 4140  
 gactgattag aagcctcgt caggggtatt tctcgcttcc agacagcaca tgactgtcat 4200  
 ttggcacgtc tttagcgtc cttccgggag aggggctgca acctggcag gcgctgtggg 4260  
 ggagggggct aggacatcct gtgctggtt tcaccaagt ggtgtgtgga ctttccctgg 4320  
 cttccccagg ctgtctggct gcacagctt ggggaaacgg cactgggtc aagcgggccg 4380  
 agaagaggaa gtctgtggtt tgtctctgct acagactggc ccagtgagg ctgtccagca 4440  
 gtgcagggca cagagcaaaa gcaggaggt atgggcctac tccccggtc gccctgtgg 4500  
 ctggctgtgg ctctgcggg tgctgacaag tactcgccc tcctgcgggt caccagggtg 4560  
 catgcccga agcctccat tcttctcctg gtttgagggt ctttctcctg caccacccc 4620  
 agcggccagt tcagctcaac ttccagaaat ctggttcacc cccaatccct ttctcataac 4680

tgcttccaag cccagacaag gagacagacc ccaaagatc cctaccccta tttcgcacc 4740  
tgaaatcgca ccacgggaag agctttgctc atagagtcaa taaggcttag agtccaggcg 4800  
cctgtgcgag ggagcaggtc atcacccttg taccacccgt ggttttagac aggaccctga 4860  
ggttgggggtg gggctggggc tggagaggag ccagggtgcc tggcccttgc ttgggccccg 4920  
tgtccctgtg atccaggctg ggcgtgctat ggggtgctggg tgatattcca gccctgcagg 4980  
tgtccgcctt gttcccagca cccctctggg caagaagaac caggctctcc cagaaatggg 5040  
cttcagtgat ctccacttcc aagtcgtccc cacctgcctt gtaggacaca gtggtacctg 5100  
gtatgctggg cagccttcca ggaacctctg gacttactca gtgtcccca gccctacaca 5160  
ccattctttg tgtttctggg cccaaactaa gcccccaac ctgggctgca gagcaagtgc 5220  
tgaatcatga gagacccttg agggtcctcc aggtaggccc ccagtgtgg aggagtcccc 5280  
tcaggcaggg ggccacgccc aagggtgtgg aaggctcagct ggcagccgga tctcactttt 5340  
ggggctgtag gcttcttgca ctggccgcca atgccatggc cgtgggatgg ccaggataag 5400  
gcatctgccc cccaccccca cccccgcac aaggctcttg agggctgcgg gctcaaggag 5460  
ttggcggtag ggctggggga ccaggggcac agagcttgta agcgctctc tccaggatgt 5520  
gggtggccca gcaggggagc tttgagagtc cagggtgtgag attccaaatg ctaggggcct 5580  
gagaggaggg agccaccagc ttggccagag cctgggtggat cacgccccca ccacgccttg 5640  
cccttctctc tggatcatgt ctctccacc acgtttggaa agttactgct tccctcttcc 5700  
tcagcccctc gggctcccag ttatggaagt ggcgtgattc agagaaggta aaggatggga 5760  
gggagagggc tgggtgatgg gggacccgc agggcgccct gtgctgttac atggagctcc 5820  
aggatcaggg cagggtggga gcctggggtc ctacttctc tcccagcca ggccaggctcc 5880  
ctcacagccc tgccaggagc atgatatccg ctgcggtgca gaactaatct caaagctcaa 5940  
accaggttaa cagtgtaggt aaaacagatg acagggcatg agactacccc caggacaggc 6000  
gaaggaccca ggccgatggg ggcccagaac agtcctgac ctggagctcc tcccagtg 6060  
ggacccaggg ggtttccgag gggcttagag tagggcttag aggcttagag tagggctagg 6120  
gacttctggt cttccctgcc tcgggaacag ctggctctgg aaggggcttg gtccctgggg 6180  
cactggtgcc caccaccct gatgcctggg agacaccagc atcctctgag catgtgtgcg 6240  
tcctctggt cccgagggaa gtgactctc acatcccca gctggcgggg ccagagggcc 6300  
agcatcctcg cctgacacct atttttagat gctgagacag gcggttcct cggggccagg 6360  
ggccctgtga gtggagcttc cgcttctgg cctaggagag aattcctgct cctcttccct 6420  
ccatgctgcc ttttcgcccc tggaggccac aacggggctca gaggggcagc tgctcaccac 6480

ctaggagggc ctgagagggc cctacgtcac ccaggaggga gtctggcccc gtccccaacc 6540  
tccacaccca ggcctggcac tgccccttct tgggtgggcag agagtgaggg gttggcctgc 6600  
agggacccag gctggagggg ccgttcacct ccggccccca gcgtcccttc ctggaagcac 6660  
cttggtgagc ccctcccctc cttcacccag tatctccagg ggtacttcct cctttccttc 6720  
ctgcctcagg gcctcactgt cctcctgggg aggggtgtctc aggccccagc acctcccagt 6780  
ggctgagccg aatgggcact tcccgggtgtg tttcccatat gtgcagtccc taggtgtcgg 6840  
tgagcaggca cagagcccg cgcgtggccc tgcctgggtg accccctccc caagagcatc 6900  
aaggaggggc ctggactaga gacacacaga tgcccagcct gtacgtaaag gcgggtgagc 6960  
tgatgtacca tegtccctgt cccccactgg ggtgcctggg caggacttgg ggtgaccact 7020  
tgccccgtct ggggtgggggt aagggtatggg tggggcgacc agatccctgc cctttcctgc 7080  
agctgtgggg gtgtgtgtgc tggcctggag agctcccacc cgaagttctg gctcctggct 7140  
gtccggggcc tgcgggggca gcgagcagct ggcattgggtg ggggagctga cctaggcctg 7200  
cccgggcagc gcctgtctgc ttttgcctcc ttccagctgc ttcttggaac cagcggacag 7260  
gctgggcagg aaccagtggt gcttggcagc ccccttttta aagtcgattc tgttatttat 7320  
taattcccag gaaggagaaa gaaagaaaca atccttcata gagtacaaac actgctttta 7380  
gtagccttgc aaggagccct ccaggaacct cacaggttac ctgggctcca tcctgagagc 7440  
cacctccat ccccaatccc cagcagagca tcttgtgggg tggggcggtc tgtggggcgg 7500  
ggcgcccttg gaggcgggggt gtctcgggaa gcggggcgctc tcgggaggtg ggggtggctt 7560  
tgggggtggg catttcctgg ggtggggcgt ctcggtgggt gggacagctt ggggggtggg 7620  
gcatctcggg aggcggggcg tcttgtgggg tagggcggtc tgtgggggtg ggcattctgt 7680  
ggggtagggc ggcttgtggg gtggggcatc ttgtgggggt ggacggcttg tgggggtggg 7740  
catctcggga ggtggggcat ctctggggcc cggccacttg ggaggcgggg catcctgggg 7800  
gcggggcatc tcagagggcg cctccggagg ctggagtatc ttgggaggtg ggagcaggtg 7860  
gcagagaggg ttcccacagg tgagctttga gcaggaggt gcctgtatgg atggctctgt 7920  
ggggagaggg gtgacaggag ttccagattc cggcacttat gaaacctcac agtgatggag 7980  
agccgagtgc tgcgtgtcag gctaagttgt gtgcattgca gcttctgcac ttttatttcc 8040  
ttgtttgtag acaaggcaga gagaagctga gatgggcctg aggtcgcctt ggtgaaaggg 8100  
actcagcagc cagggccttg ggctgcctc cctcatcacc gtgaaagcgg gactctcttt 8160  
taactgacat cgggctccat agttactcca gtcctaactt tgatggatcc taaaagtgca 8220



cttctaagga cgcgggttcg gtgttttccca tgccgctgct tgcccctggg aagcgttggc 8280  
 tctgcctcgg aagaagttag cgccaagatg gcagcctggg gtcttttgggg ccagaagaa 8340  
 aactggccc cggggagttc agtcatcagg gacttaggat gtgggggctt ttcaaacagc 8400  
 tttatttaga cgtgattgac acacagtaaa tacagatgtt taagggtaca acttggttaag 8460  
 ttttgacaaa tttatacccc cgtgaaacca tcaccaactc cccaggtgcc cctggggccc 8520  
 ttgggatctc tgcttctcgc ccctcctccc cgtcccaggg caaccacggg cgtcgtctgt 8580  
 ggggtgacac agcatgcatt tcttcaacaa gcggactcag aaggcacttg cacatcgttg 8640  
 ctgttctgcc tctttgcttc agcatgatta cccagaggcg cacccggtgc gtggcctgcc 8700  
 cgtcgtctat gcaccogtgc tgtggcgtgc cgtcgtctgt tgtggcatgc ctgtctgtgc 8760  
 acccggtctg tggcgtgccc gtcgtctgtg tggcatgcct gtctgtgcac ccgtgctgtg 8820  
 gcgtgcccgt cgtctgtgca cccgtgctgt ggtgtgcccg tcgtctgtgc acccggtccg 8880  
 tggcgtgccc gtcgtctgtg caccogtctgt gtgggtgtgc ctctgtctgt tctttttatt 8940  
 gccgggcagg gttgcacca catgtgcaag ccagcgacgg accccagggt caccogttca 9000  
 ccggtcagtg ggcataatggg ttgtttcagt ttggggcatt tacaagaaac gtgctagaac 9060  
 atttgtgtac aagtcttgtg tgaacctaaag ttcatttctc ttgggttaa atctgtgcgt 9120  
 ggagcagctg ggtcatgtgg tgaatgtggg tttactgct taagcagcag ttttacataa 9180  
 ctgcaaaact gttattcaag gtggctggac cgttttacag ccccggtgt atgcgtcca 9240  
 gttgcctccc ccagcagcat gtgggtgtgt tgggtctttt cgtggcagcc agtccactgg 9300  
 gtgcgtcgg catgtggctg cagcttgacc tgggtttcct ggtccctggc aagggtggagc 9360  
 atctcttcat gtgtttttt gctgtgtgtg gatcttgccg ggaagggtct gttcctgttt 9420  
 tttgcccac tttcaaagat tgggttgcca gttttcttgc tgttgagttt ggaaagctct 9480  
 gcatacgttc agggcacagg tcttttacca ggctctgccc caggctcttc ggagagcagg 9540  
 tgtctttcgc attcctgact ctggggaacc tctagccctg ccacatgggg tttgttatgg 9600  
 ggcaggggca cctgtgcctt tcccaccacg gggcttgggg atttggtgct gccattgccc 9660  
 tccctcgtag gtggccctag gggggtcct ccgctccgt ttctcatcc agaaaccggc 9720  
 agtgaccatc accaccattg ttgtcaccta gctccagctc aaggctccctg ctgaaggctg 9780  
 gagagcttgg catggccccg tttgtccatg ctagggtggt gaagaccaag gctcagggtga 9840  
 ggctctgcc cagtgcctgg cactccttct tgccccattt ttccaccag ggtggctccc 9900  
 gactacttct ggtagcctcg gggacagttg aggtggacag gctggcgtca cccccattc 9960  
 cggctgtccc tcccacccc tctgggcca gctgttctgc cctattaaaa gtcacatggg 10020

ccctcgggtc cttcctggtg ttggcccagg ctctttcagg ccctgcaggc caggaccagc 10080  
cttccttgca accctcggca gaggcctggg gccggggctt gtctaggggc agcctcccca 10140  
tacggccctg gagtctgaac agaagcccct tcccagagca cagcaagaag ctgcaacgtg 10200  
gcctgaagtc ccaccattag cagggtttggg gtttaggctg agctttgcca tcactacctt 10260  
tctgttagga cggtatgccc attagatggg atcatcccct cagcgcccag gctagaggag 10320  
gggtggtccc tgcccagcca gggagggctg ggggtggatg ggcctctaca gagcagcttc 10380  
cgagccaggc acggttccat gatcagctct gttttataga gggggacact gaggaaccgg 10440  
gagcctgggg accttccagt ggccccacag ctccctgtggc tgagtcaggg tttgtcacca 10500  
ggcctctgtg gggatgaggc tccccatcc acctgcccc aacctgtcctg gaacagctct 10560  
caaaacggtc tctggaccac agtttcaaaa gaaaataagc aatgttttca aaggccctgg 10620  
aggaagccag agttaccag gcaactctcg gcctcgccac ctccctccgc caggctgcat 10680  
ctggagccag ctcaggaggg cagcagggtg aggacagcca ggctctcttg ggccaccccc 10740  
cagccccac ctttctgccc tctcctgcac tgtccacggc cctccctgtg ctccacggg 10800  
tataatgggc acagaagaac caggagctgt ctgcccctgc aggattcttg aagccagggg 10860  
cccctggcct ccctggggcc ttgtcatgtg aggggcacac gtggggctcc agctgccaca 10920  
tggtctccag cgctgcccgc aggtgtatgt tggggccttg gtgactctaa tgcaccttcc 10980  
actcggcaca gaagagcttc agtctggggc ctggggcggg gaagtaggct gccatcctcg 11040  
ctaaaccaa gtgtgaaaat tgagttgaaa ctcccatagg agggcaggag gcacagctcc 11100  
tcagaagaag gtctgagaaa ccacagccca gggtgtgtgt tcgggtgtgt ggagaagggtg 11160  
ctctggcagt cctgctacag ggggaccatc aacagcccct ttgggggtgag agccccgtg 11220  
ctgctggcac cagcagcccc tatgaggctt attttatttt tgagacaggg tcttgcctctg 11280  
tcaccgaggc tggagtgcag tggcacaatc ataactcact gtagcctcaa cctcctgagc 11340  
tcaagcgatc ctctgcctc agcctccaaa ggtgctggga ttacaggcgc ttgctaccac 11400  
gcccagcccc ctctggcctt attgtttgcc agggccagct caggctcccg aggaggggag 11460  
acaggagtgt gagggaaagg gggaagaggt atagagcccc cagctcctcc acccaccoga 11520  
accctcaccg aggcctaga ccctagaccg gcctgaccgg ggggtcctca ggccggggac 11580  
ttgggtgcag gccatggtgc tggggcctga agctcacgct ctgctgagca cagccccctg 11640  
cccaaccca ccctggggcc ctgcttccct ggccagggcc attggaacag gagtggggct 11700  
gtccaggtgg tgttcttggg tccagccctc agtttctctt ctgcagttga ccggcagccc 11760

tgcatctgtg gtgggggtcgg cgcctgggtgc tggtagaggca aggcctcagc tgctgggaca 11820  
ggacctgcct ggcaccacagc tggtaggcaga gccaagcatt ccgactcagc tctgggagca 11880  
gctgccttct gggctggcat tctccgccag ggggggttggt ccctcgtggc cccccccggg 11940  
tgccctccta cctggctgat ttcatctcct gtccccctgc ctctcctcc aggaagcccc 12000  
cagggcctgg ccctccttga gagtggcatg gaggaggaag aagactcgcc cagggccatg 12060  
ggagtcggat ggtggccgca cttgtggggc cctgacccca taggcttctt cagcacgccc 12120  
tggcctgggt gatccctgcc tgagggtgtg gcacggctca tctgccagac cagatttttag 12180  
gggattcttg tactgtcctc ctggagcagc aggggggtaaa gcctgaccca ccagactgt 12240  
ccagcaacaa gggcctcctg ctgtgggcca gggaccctgg aactgaccaa ttgtgtccta 12300  
gggacgcaga gtccccaggc tgctagaggg ctgtggggcc ctgtttcatg cctgaagcag 12360  
gaagaaaccc caggagaggt ctgaaggga cccagcccc accctgtcta gcagggagga 12420  
gcctctgcaa gaggccgagg ggtgctgaag tggaggagga tagaggcagc aggactcagg 12480  
gtcactggtc atttatggg atcacacggc tgcaagtgtgc cctgcatggt gctaggcacc 12540  
agggacagca gaggacaagc ctgtgtcctc tcccaccacc agagggtggt gcactgcccc 12600  
tagggagaga gggggccttg gtgtgtgcag aggggggcct ggggcacgtg cctggcctgg 12660  
tcagatgatc agagtgggt gggctgggccc tgggtctggg cccagtctca agggcagacc 12720  
ccacctggct agagttgatt gtgtgcacac cggatgacct ggcgttgaag gcctctcctc 12780  
tctgtgagcc tcatccccac ctgccagact cccagcacag cctgcttctt gccccagctg 12840  
ctgagcgaca gcgctgggccc ggcttctgcg cgccttctcc cccagcccat cttggaaacc 12900  
acagcagcgt ctttctctcc aagtcccttc ccagggtgta catccacag cagggatgta 12960  
tcccacaaac cccgaggccc ctggtgccta cagcttggcc tggtaacatc aaatcctacc 13020  
ctctcctcct ggcagcaaag atggggtgcc cccaccccag agttctcagc acccccagac 13080  
agaagcagtc cccagcgac ctcagaactc ttggggcgct gccacacct tgcaggaggg 13140  
ggcagtgttc ctgggatgct caggctcctgg tatcacctct ggccagatac ggaaggtgaa 13200  
actacagggc atccaattca ccttgaactt cagataaaca ccagattatt tttttgtatg 13260  
tcccgtgcaa tatttgggac acacttacct taaagaagta ttctgttttc atctgagagg 13320  
cagatttaac cggcgtcccg tgtcttctctg gcagtcctgc cctggagtca cactccacag 13380  
gtgcagggca gggccaggct ccaagtagat ggcggccaaa gcacccgccc catgctcctg 13440  
actcccgggg ctcttcaggg cattgcgaaa accagcagca gagctgacac ctgggtccctg 13500  
ctcgggagcc agcaaggcag gaggtgctt aggccttgcg tgtgggggtg gcgcactccc 13560

tgtgtcagtg ctcttcgtac atgtgacaact gttcccgcgc tttcccagct ggctggaggc 13620  
gtgatcctgg gtgtggccct gtggctccgc catgaccgcg agaccaccaa cctcctgtat 13680  
ctggagctgg gagacaagcc cgcgccaac accttctatg taggtgagtg cacatgtggc 13740  
cgcagacgca ttcagggagg gcttctagga ggaggcaggt cctagccttt tggatgggga 13800  
catggagggt gaaagacagt cgggcatggc gtgtccgggc agggaggcgg ccctggaaag 13860  
ggctctgggc acaagggttg agatggagggt gggcctgtgg cctgctggcc cttctggtct 13920  
gagccagggc aggggggtggc agctaggcct gggcagggac tgtgtggaga ccttgcttat 13980  
tttaagtgtg gggttatttc gggggaggct ccctgagaag ggtggggctg gatgcctggg 14040  
ccacacagag cagccgaggc agctggcgct gtggagcccg ggaggaggag agggatggag 14100  
ctcaagggat ggaaccagt gaggggtgga gacggggcag gggagggggtg gagaggggtg 14160  
gagacgcccc agaggcgggtg tgactcagct gccctgcag gcagctgcac cttgctgcct 14220  
tattaggctg cgtgtggggg actgggctgc cctccctgcc cccaggagca ggagcaggag 14280  
tgatggagga ggaggaggag aggggcaagg ccaggaggag gaggagggcc atctcactgt 14340  
gcagagagca gcacccttc tcctgggtgcc cctggcaggg ctggtgctgg tggggctctg 14400  
ggagcatttg ttgagatgct tctggccttg aaaggaggcc cctgggatgg ctctgttgcc 14460  
ctcacaggct gaggggtggg tgagggtggc agcctgtgtg tcccagtc tcagggcttc 14520  
cctcagccgg caggtgcccc caggcctgga gctgcagggc caggccccct gccagttacg 14580  
gaggctgctt ggcttggttg ctgaaccagg gcccaggag gccgaaatag cccacacct 14640  
gogccgtccc acctctttgt ccagtcaccc caggggcagg tgagggccct ggccacacag 14700  
cgtgcccggt ccttcttccc catgccccgc tcatgggtca gagggccggt gctggggctc 14760  
agatggtgtc aacagggatg gtccctgtcc tcccagaga cagaagcctg tggcccacgg 14820  
agggtttctg ggcccagccg atcctaggga ggggtccatg gccctgccc taggttcctg 14880  
gcctctctcg gggccgtggt gccctcacag gtggtgtcag gaaggacggg aaaggctgct 14940  
tgtcccaggg gctcatgtgg agaccacccc ctgcaogcag ctggggcgct cctgcctgtg 15000  
tcctcagaag cactcggtt agctttgcc atgtgcctgg gctgtgggtg gcagagccc 15060  
gccagcatcc tccgatctcc aagggtgcat ctctactgga ggccctcct gggcctcttg 15120  
ctccccgctt cccagatcat taggatattt ggggtccaga agggcctccc agccatcctg 15180  
ggccttgctc tccggggcca ccagtccagc cagtgacaac cacagcatcc ccggcctgga 15240  
acgaggctgc cccagcacg ttcctcgtag tcctgtccag ggacaggagg ggctgcccct 15300

gccaccgagt ccccttctcc aggacctggg gcctgtgggt gtgaggcagg tgttcttga 15360  
aggggtcact ctccaggcac ccggcggcca aggccttgtg ctggagcagc tcccgtgtg 15420  
gggtcggcgt cgggccccgt gtggccggag aggagctgaa gggtcactta gcttcgggct 15480  
ggggcgagga caggggacac ccagagagg tatgccaggc ctcttcctg cgcacctc 15540  
tcggcagaag cagaggtcac aggcctgtgt gaggcccat ggtgctgcc ccgatgacc 15600  
agggtgaggc tggcgttga agcagggtgt tgacctgcat ggtgtaccg tggccacatc 15660  
agagctccag cccagagcc gccaccctc ggtccttggc tgtggtttcc ctgggctgga 15720  
ggagcctgcc gttgtgttg ccacacgacc acaggacctg ccacccccga cgtgggctct 15780  
gcctggggcc cactggaca gggaccctt ggagctctc tggccacaa gtcctcgccc 15840  
attccagaat cggccttctg gagcctcttg ctgtccctga tgcgggctgg gccttgccaa 15900  
gggctttttt tctgcgccc ggaacagggt ggatttctg ggctcactcc cctcagagac 15960  
gctgcgggtg cgggtgggtta ggcccaaggc cgttaagaga ggaggctggg gtgggctgg 16020  
ggcctggcag ggggtctggc agccctgggc ctccacctc ctgtcaggac caaaaaaggc 16080  
aacgcgcctc tctgacctg taccocggag tgaaccaac cttgcaacct aggagtgtca 16140  
gggcctgagg ggaggagac ctggctcctg ggtgccgtgc ccgtaaggag gtggccacct 16200  
gcagggcatt cctggcagag gcttcatctg gccaggtagg aggcctgggtg gccgagcccc 16260  
aaatctgggt gtgttctctg cctggcggtg ggtcctgcc caggcacctt ctctctggg 16320  
ctggctgggc agggacaatg ggccctggct cgaggagggg gcctgggctg cttctgcat 16380  
tgctcgggtg acgggagatg gccctgcct gctgagggat aggggagtgg gcaggcagt 16440  
agagacactg acagctgtcc cgggggtaca gggccctgtc tgggtggcca ggcccatgtc 16500  
tcgggcccac agtgcgcccc ccaccttgg acggcgctt ctccctccc aggtgcatgc 16560  
tgcccagcca gggagcgtg gggagtctg gagggtggc ctacacgccc tggccagct 16620  
gtcccagggt ggggtgctgg cttcagccct cagcccagg cctaggaatc caacttgatc 16680  
ctccccacac agcagccagg ttcaaagtca ggtcccgtaa cggaagtgt gctgtgcagc 16740  
ccagattggg gggcaggagc cagcagggcc cccccacct cttctcgac cactgagg 16800  
aggcagcatt ggttccagtt ccggttctg ggtgcctc tcaaccccg cctacagtgg 16860  
ggccacct gtgccttctg atgccactcc caccacgc caagtcccag aggccttggg 16920  
agcgggtgaa ggcgggtggg ggcgggtggc aggtgcaggc ggtgggtggg ggggtggca 16980  
ggtggcgggc cccaccgag gtgtcatccc tgcgaagcac ctgtcgccag cactcagagc 17040  
gctcatgagg tgccagtc ccagtggcc tcttagtct ccgtcctgtg tcatggaaga 17100

ggtaactgag gcacagaaaa ctaccaggc caggctggga tgtgaggtcc cttgctgctc 17160  
atccctggca gtcagcaacc ctacatcttc ccagctgggc ggcccgtggt gggttcggca 17220  
cccaggaccc tccgggtct tgggctgtgg cgagtgtgta ggcaccacc tgggtgtctct 17280  
ctccccgcaa ggcattctaca tctcatcgc tgtgggcgct gtcattgatgt tcgttggtt 17340  
cctgggctgc tacggggcca tccaggaatc ccagtgcctg ctggggacgg taaggcagg 17400  
aggcgggcct gtgcctgggc cggggagggg ctgggggctg cgtctggccc tgaggaggg 17460  
gcagagctgg tgctcagggc ggagcctaga attctggggg aggtggctcc tgtgccctgc 17520  
ttttcccggt tgggttttaa attaaatccc accgtgcttg gtctccatcg tggccagttc 17580  
ctacgtgacc gcttttcttt gtcaaaaaat agccacaaat ataacaggga gcaagcctca 17640  
gctctgaggc cagcctcggc gtcccgggca caccgcccc tgtgggaagc ccaggcctgg 17700  
ctgtgccatc cagggcctgg ccagtcagg aagagggagc ctatgcccggt gtctccagt 17760  
ggggaaactg aggcagatcc catggctccc ccttcctggt ggagcaggaa caagggggtg 17820  
gggaagatca gtcaggggtc atgctgctgc acacgcctcc ctgggggctg cagacatcct 17880  
ggactacca gcctgtgacc ccaaaccaca cgccccgcc catccacccc gtcctgtgga 17940  
gcctggtgcc gcgtggggac atcctgggct ttgacggctc ctccctgcgc tgagttttag 18000  
cctctgtgcc ccagggtcc acacaagccg ctactcctg gtcaggctgt gggctggtgg 18060  
ctccactag cccctcacag acacgcctgc tgggcacctg ggtgtgtgtc cttgggcccc 18120  
gcctacagcc tgccctcttt cctccctctg gccactgcc ggctccagtt cttcacctgc 18180  
ctggtcatcc tgtttgctg tgagggtggc gccggcatct ggggctttgt caacaaggac 18240  
cagggtgagc tgggtgtgca gggacagggt ggggtgggtg acgggggcac cctcctctcc 18300  
tgtcgcggtt gggggttggg ctgactcatg gcttgtggga gctctttggg ctcttcctgg 18360  
gtcccacttg ccaggaggat ctccaggggc tttatggagg aggcagcatt ggggctgagc 18420  
accaggccag cctcccgtgt ccagcactc ccggggcagc tgagagtga gagtccttgt 18480  
cctctggggt ctagcctcga agccaccctg ccaggggaga gcctgggaaa agtgcgtccg 18540  
cctggggcgg ggcggggtgg gggcaaggag ggggaggttc cccctgtgca tgtgaccgca 18600  
cccctcccc agatcgccaa ggatgtgaag cagttctatg accaggccct acagcaggcc 18660  
gtggtggatg atgacgcaa caacgccaag gctgtggtga agaccttcca cgagacggtg 18720  
cgccccggg gggcgagggc ggggagcagg gccccggaa cccggcggg tgtgtctcgt 18780  
cctggatgaa tcctgcctac gccagacct caggagcagg aggtgccctt gggacctcca 18840

ggacccctgg tctcaactgg tcctcggttg ggaacctagt gggccagggt ggcccagggt 18900  
gcggaagct ctgagcagcg cagctgagga ggaagaaggc tggcccctgg atgcattctg 18960  
cagtggggag cgctgcgtac ccctggccac ctccccatgg gttccctaga gccaccgtcc 19020  
ccctgggcac atccagggt gaccttgac ccctgctctc tgcagcttga ctgctgtggc 19080  
tccagcacac tgactgcttt gaccacctca gtgctcaaga acaatttgtg tccctcgggc 19140  
agcaacatca tcagcaacct cttcaagggt cgcgaggccg gtggggccgc gcctgacccc 19200  
ccgcatgtcc cgcccctggg tggggtccta ggggtgggca ggtcacacgg cagccccaca 19260  
gggagcgacc aactgggtg gcattggccc tgtcagggt gctctgctgg gaggggtggg 19320  
gtgggaccgc atctggcca cgaggaaggc aggcgccttg tgctgcgcat tccgggtgaa 19380  
gaaggtggag gctctggggg gtgggaactc acctgcaccc ccagctccac gtgtgcactc 19440  
gtgggtgtgg acgcccctga cagcctgtag ctggcagggc ctgcaggcca tatagtgccc 19500  
tgtggaagtt tcctgctgag gcctcagtgg aagtcgtcat cagtgatgct ttaggggtct 19560  
agtgcacca atgaccgtga tctcagtga aaagggcaca gtgtgtccca ggcatttcgc 19620  
gtttatgtta aaacgggtgg aagatagcaa gccggcagag gccgggcccgc tgcacccgcc 19680  
tgttccgagg tgggtagggg gtggggggct gttcccagga ttcccctcta cgctttctgt 19740  
ggtgaccacg gattactgcg tgacaacggg aagccgggag ccgaggcccgc gtccctgacc 19800  
acgcgtgcct ggccaccct gcaggaggac tgccaccaga agatcgatga cctcttctcc 19860  
gggaagctgt acctcatcgg cattgctgcc atcgtggctg ctgtgatcat ggtgagcggg 19920  
cgggggcccga gggcctgctc tctgggtgc cccttcgcg gggccttgtg ctgactgcgc 19980  
ccccaccac cctcctgcag atcttcgaga tgatcctgag catgggtgctg tgctgtggca 20040  
tccggaacag ctccgtgtac tgaggccccg cagctctggc cacagggacc tctgcagtgc 20100  
cccctaagtg acccgacac ttccgaggg gccatcaccc cctgtgtata taacgtttcc 20160  
ggtattactc tgctacacgt agccttttta cttttggggg tttgtttttg ttctgaactt 20220  
tcctgttacc ttttcagggc tgacgtcaca tgtagggtggc gtgtatgagt ggagacgggc 20280  
ctgggtcttg gggactggag ggcaggggtc cttctgccct ggggtcccag ggtgctctgc 20340  
ctgctcagcc aggcctctcc tgggagccac tcgccagag actcagcttg gccaaacttg 20400  
ggggctgtgt ccaccagcc cgccgtcct gtgggtgca cagctcacct tgttccctcc 20460  
tgccccggtt cgagagccga gtctgtgggc actctctgcc ttcattgcacc tgtcctttct 20520  
aacacgtgc cttcaactgt aatcacaaca tcctgactcc gtcatttaaat aaagaaggaa 20580  
catcaggcat gctaccaggc ctgtgcagtc cctcagtgcc agtgggtgtct gagacctagg 20640

gggtggccgg agggcagggg aatctgacat cggtggggct tggctctgtg gactctgtgg 20700  
 ggtccagggg gaggggtgggt gggtcgggat ccctgggtgtt caccaaagga gtcactctgt 20760  
 aaaatttggg gagttattta ttctgagcca aatatgagca ccggtggcct gtgacacagc 20820  
 ccaggtcct gagaacttgt gcccaaggcg gtctggctac ttaattgtat acattttagg 20880  
 gacataggac attgatcatt acatctaaga tgtacgttgg tttagtcgga aaggtgggac 20940  
 gatttgaagg ggagggactt tcaggtcata ggcggattaa aagatgttct gattaataat 21000  
 tggttgattt tatctaaaga cctgaaatca atagaatgga ctatctgggt taagaggagt 21060  
 tgtggagacc aagattatta tgcagatgaa gccgccagat tgtaaagtgt tcttatcaga 21120  
 cttaaaaagg taccagaatc ttagttaatt ctctcctgga tcaggaaata gacctggaaa 21180  
 gggaggggga ttctctatag aatgtagatt ttcccaagag acagctttgc agggccattt 21240  
 caaaatacat cagagaaata tattttgggg taaaatactt cggtttcttt cagggcctgc 21300  
 tgtcacgttg gtatcttatt actacagagt ctgttttgtg agtcttaagg tctttttatt 21360  
 tttagacaga gttttgctct tgtcaccag gttggagtgc aatggcgtga tctcagctca 21420  
 ctgcagcctc ccctccacct ccaggttca agcgattctc ctgcctcagc ctctgagta 21480  
 gctgggacaa caggcatgca c 21501

<210> 12  
 <211> 000  
 <212> DNA  
 <213> Homo sapiens

<220>

<400> 12  
 000

<210> 13  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 13  
 gcgcctttcc agcacaatgg

20

<210> 14  
 <211> 20  
 <212> DNA



<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 14

agccagccag aagacgaaat

20

<210> 15

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 15

cctggctgag caggcagagc

20

<210> 16

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 16

gcacttggtg cagccctcca

20

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 17

ttgaagacga agagcaggta

20

<210> 18

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 18

21

cgaaattgaa gacgaagagc

20

&lt;210&gt; 19

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 19

gccagaagac gaaattgaag

20

&lt;210&gt; 20

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 20

cctccagcca gccagaagac

20

&lt;210&gt; 21

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 21

ggatcacgcc tccagccagc

20

&lt;210&gt; 22

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 22

cacagggcca cacccaggat

20

&lt;210&gt; 23

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23

cttgtctccc agctccagat

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 24

gcctacatag aaggtgttgg

20

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 25

atgtagatgc ctacatagaa

20

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 26

gatggccccg tagcagccca

20

<210> 27

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 27

gtccccagca ggcactggga

20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 28  
acctcacagg caaacaggat

20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 29  
gttgacaaaag ccccagatgc

20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 30  
acatccttgg cgatctggtc

20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 31  
gcttcacatc cttggcgatc

20

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 32

gaactgcttc acatccttgg

20

&lt;210&gt; 33

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 33

tcatagaact gcttcacatc

20

&lt;210&gt; 34

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 34

cctggtcata gaactgcttc

20

&lt;210&gt; 35

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 35

gtcttcacca cagccttggc

20

&lt;210&gt; 36

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 36

cgtctcgtgg aaggtcttca

20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 37  
gggacacaaa ttgttcttga

20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 38  
gcccgaggga cacaaattgt

20

<210> 39  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 39  
ccttgaagag gttgctgatg

20

<210> 40  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 40  
ggcagtcctc cttgaagagg

20

<210> 41  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 41

aggtacagct tcccggagaa

20

<210> 42

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 42

ccgatgaggt acagcttccc

20

<210> 43

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 43

caatgccgat gaggtacagc

20

<210> 44

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 44

atggcagcaa tgccgatgag

20

<210> 45

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 45

gcgaccacga tggcagcaat

20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 46  
gatcacagcg accacgatgg

20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 47  
aggatcatct cgaagatcat

20

<210> 48  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 48  
agcaccatgc tcaggatcat

20

<210> 49  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 49  
cagcacagca ccatgctcag

20

<210> 50  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>



<223> Antisense Oligonucleotide

<400> 50

tgccacagca cagcaccatg

20

<210> 51

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 51

ccggatgcca cagcacagca

20

<210> 52

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 52

ctgttccgga tgccacagca

20

<210> 53

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 53

cggagctgtt ccggatgcca

20

<210> 54

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 54

gtacacggag ctgttccgga

20

<210> 55

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 55  
gggcctcagt acacggagct

20

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 56  
gcagaggtcc ctgtggccag

20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 57  
tacacaggcg gtgatggccc

20

<210> 58  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 58  
tacgtgtagc agagtaatac

20

<210> 59  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 59  
aagtaaaaaag gctacgtgta 20

<210> 60  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 60  
aggtaacagg aaagttcaga 20

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 61  
catcagccct gaaaaggtaa 20

<210> 62  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 62  
ctcatacacg ccacctacat 20

<210> 63  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 63  
gcagaaggac ccctgccctc 20

<210> 64  
<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 64

gagcaccctg ggaccccagg

20

<210> 65

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 65

aggcagagca ccctgggacc

20

<210> 66

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 66

gcgagtggct cccaggagag

20

<210> 67

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 67

ggccaagctg agtctctggg

20

<210> 68

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 68  
gtgcagccca caggacgggc 20

<210> 69  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 69  
gaggaacaa ggtgagctgt 20

<210> 70  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 70  
cacagactcg gctctcgaac 20

<210> 71  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 71  
aggcagagag tgcccacaga 20

<210> 72  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 72  
gcatgaaggc agagagtgcc 20

<210> 73  
<211> 20  
<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 73

gtgttagaaa ggacaggtgc

20

<210> 74

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 74

tacagttgaa ggcgacgtgt

20

<210> 75

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 75

tgtgattaca gttgaaggcg

20

<210> 76

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 76

acggagtcag gatgttgtga

20

<210> 77

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 77

ttattaaatg acggagtcag

20

<210> 78

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 78

tccttccttta ttaaagtcag

20

<210> 79

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 79

agcgacacca gtggactggc

20

<210> 80

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 80

ccaggcctct gccgaggggtt

20

<210> 81

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 81

ggcctcagca cagcctgtga

20

<210> 82

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 82

acctcccca gaattctagg

20

<210> 83

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 83

aggtgaagaa ctggagccgg

20

<210> 84

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 84

ccaggctcac ctggtccttg

20

<210> 85

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 85

agcagtcaag ctgcagagag

20

<210> 86

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 86

gcccgctcac catgatcaca

20



<210> 87  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 87  
ctggctggtg gccggcaaag

20

<210> 88  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 88  
gtaacaggaa agttcagaac

20

<210> 89  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 89  
tcagccctga aaaggtaaca

20

<210> 90  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 90  
tttttgactg cacaggcctg

20